

5%

Sevin® Dust

N-Methyl Carbamate

ACTIVE INGREDIENT:	BY WEIGHT
Carbaryl (1-naphthyl-N-methylcarbamate)	5% by wt.
INERT INGREDIENTS	95% by wt.
TOTAL	100.0%

EPA Reg. No. 10107-43 EPA Reg. No. TD-07-NE-1

For MEDICAL and TRANSPORTATION Emergencies ONLY, Call
Chemline 1-800-424-9300

SEVIN is the registered trademark of Rhône-Poulenc Company for Carbaryl insecticide.

KEEP OUT OF THE REACH OF CHILDREN**CAUTION****STATEMENT OF PRACTICAL TREATMENT**

GENERAL: Contact a physician immediately in all cases of suspected poisoning. Transport to a physician or hospital immediately and SHOW A COPY OF THIS LABEL TO THE PHYSICIAN. If poisoning is suspected in animals, contact a veterinarian.

IF SWALLOWED: Never give anything by mouth to an unconscious or convulsing person. If conscious and not convulsing, drink 1 to 2 glasses of water and induce vomiting by touching the back of the throat with finger. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

SEE ADDITIONAL PRECAUTIONARY STATEMENTS INSIDE BOOKLET.

Manufactured for
Omnibit Chemical Company, • P.O. Box 410 • McCook, Nebraska 69001
2001

BROCCOLI (3-Day Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Flea Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

PRUSSIAN SPROUTS (3-Day Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Flea Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

CABBAGE (3-Day Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Flea Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

ONIONS (No Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 10 to 20 lbs. per acre. Spined Leafhopper — 20 to 30 lbs. per acre. Lygus Bug, Spittlebug, Cicada Egg — > 20 to 40 lbs. per acre.

CULTIVATED (3-Day Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Flea Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

CHERRIES (1-Day Pre-Harvest Interval) — Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Twospot Moth — 20 to 100 lbs. per acre. For Codling Moth, apply at 100 lbs. and every 10 to 14 days thereafter until control is achieved. Black Cherry Aphid, Brown Salt Scale, Cherry Fruit Fly, Cherry Fruitworm, European Bud Moth, Forest Scale, Plum-Twig Leafroller, Japanese Beetle, Larval Scale, Mossy Plum Aphid, Oyster Shell Scale, Peach Twig Borer, Plum Curculio, Prune Leafhopper, Redbands-Twigroller, San Jose Scale — 30 to 100 lbs. per acre. For plum curculio, apply when curculios are present. For Plum Curculio, apply at 100 lbs. and every 10 to 14 days thereafter until control is achieved.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoe plus socks
- Chemical resistant headgear for masks and facegaurd

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions or warnings, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use dust control systems, and/or jobs, or a task in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (c) (1-8)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, chewing, chewing gum, using tobacco, or using the toilet
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

MAY BE HARMFUL IF SWALLOWED. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes or skin or clothing. Wash thoroughly after handling.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Acute is anticholinergic. AVOID CONTAMINATION OF FOOD, FEED, FEEDING THROUGHS, AND WATERING APPARATUS. KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. STRICTLY OBSERVE LABEL DIRECTIONS AND CAUTIONS, AND APPLICABLE FEDERAL AND STATE REGULATIONS. DO NOT USE ON ANY CROP NOT LISTED ON THIS LABEL OR SUPPLEMENTAL LABELING AS ANY USES NOT LISTED MAY BE ILLEGAL AND/OR HARMFUL.

1

CHURNED CABBAGE (14-Day Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Flea Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

CLOVER (No Pre-Harvest Interval) — Flea Beetle, Mexican Bean Beetle — 10 to 20 lbs. per acre. Grasshopper — 10 to 30 lbs. per acre. Use 20 to 30 lbs. rate controlling nymph. Repeat as needed. Alfalfa Caterpillar, Cucumber Beetle, Green Cloverworm, Japanese Beetle, Leafhopper, Thrip, Commerical Alfalfa Hopper, Tarnia, Vankampen Caterpillar — 20 lbs. per acre. Army Worm, Corn Earworm, Flea Bug, Webworm — 10 to 30 lbs. per acre.

COLLARDS (3-Day Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Flea Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Collard leaves after edible parts start to form, apply at 5 to 7 day intervals.

COCONUT (No Pre-Harvest Interval) — Aphid, Coccus Sawfly (if a coastal region of coast during the pollination period with a steady sawfly population), Corn Earworm, Corn Rootworm adult, European Corn Borer, Fall Armyworm, Flea Beetle, Japanese Beetle, Leafhopper, Salt Scale — 20 to 40 lbs. per acre. For larvae in the whorl (budworm) damage and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 4 day intervals starting when silks first appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestations. Timing and good coverage are absolutely essential for effective control.

COTTON (7-Day Pre-Harvest Interval) — Aphids, Cotton Fleahopper, Cotton Leafworm, Flea Beetle, Spotted Salt Scale, Thrips — 10 to 20 lbs. per acre. For early season control of cotton Leafhopper, Flea Beetle and Thrips. Claims for aphids are limited to population suppression when used with a repeated application. For concentrate dust, calibrate ground equipment or aerial apparatus accurately. After properly adjusting equipment to deliver the required low volume, maintain constant speed throughout application to insure even coverage. Strictly avoid excessive application. To effectively apply concentrate dust

2

WALU (14-Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

KOHLFABT (3-Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

LETTUCE (3 Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

LOGANBERRIES (7-Day Pre-Harvest Interval) — Japanese Beetle, Leafhopper, Larva Roller, Raspberry Aphid — 40 lbs. per acre. Climbing Curculio, Orange Tortrix, Sawfly, Shiny Tree Curculio — 40 lbs. per acre.

MELONS (including Cantaloupe, Honeydew, Honeyball, Muskmelon, Persian Melon, Watermelon) (No Pre-Harvest Interval) — Sevin injures watermelons in Florida. Fleaworm, Melonworm — 10 lbs. per acre. Injures watermelons in Florida. Cucumber Beetle, Flea Beetle, Leafhopper, Squash Bug — 20 lbs. per acre.

MUSTARD GREENS (4-Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

NECTARINE (3-Day Pre-Harvest Interval) — Codling Moth, Fruit Tree Leafroller, Japanese Beetle, Orange Tortrix, Oriental Fruit Moth, Peach Twig Borer, Persimmon Moth, Tussack Moth — 20 to 120 lbs. per acre. For Codling Moth, Persimmon Moth, Tussack Moth, apply at 50 lbs. per acre. For Orange Tortrix and Oriental Fruit Moth, apply at 50 lbs. per acre. 10-14 day intervals. Full control is achieved.

2

CKEA (No Pre-Harvest Interval) — Corn Earworm, Stick Bug — 20 to 40 lbs. per acre.

PARSLEY (3-Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

PASTURE AND RANGE LAND (No Pre-Harvest Interval) — Refer also to GRASS, Mosquitoes (adults) — 5 to 15 lbs. per acre. For Cattle cluster application only. Thorough coverage is necessary for effective control. For optimum control, apply in early morning or evening when air is calm and adults are active. 5 to 20 lbs. per acre for low pressure hydrant or ground cluster spray. Grasshopper — 10 to 30 lbs. per acre. Use 10 to 20 lbs. on sparse vegetation and for controlling Nymphs. Armyworm, Stick Bug, Thrip — 20 to 30 lbs. per acre.

PEACHES (1-Day Pre-Harvest Interval) — Caterpillars, Oriental Fruit Moth, Peach Twig Borer, Penncross, Flea Beetle, Plum Curculio — 20 to 200 lbs. per acre. For Oriental Fruit Moth and Plum Curculio apply insecticide for 10 days to achieve satisfactory control is achieved. A minimum of 10 days of control is required to insure adequate coverage. Redbanded Leafroller — 40 lbs. per acre. Codling Moth, Feltus Scale, Fruit Tree Leafroller, Japanese Beetle, Persian Scale, Olive Scale, Orange Tortrix, San Jose Scale, Tussack Moth — 50 to 120 lbs. per acre. For Codling Moth spray at petal fall and every 10 to 14 days. Inadequate control is achieved.

PEANUTS (No Pre-Harvest Interval) — Blister Beetle — 10 to 20 lbs. per acre. Grasshopper — 10 to 30 lbs. per acre. Use 10 to 20 lbs. spray on young plants and for adult Diptera, Adults Caterpillar, Cicada, Cowpea Curculio, Japanese Beetle, Leafhopper, Three-Spotted Leafhopper, Tree Vole, Webworm Caterpillar — 20 lbs. per acre. Armyworm, Corn Earworm, Stick Bug, Webworm — 20 to 30 lbs. per acre.

PEARS (1-Day Pre-Harvest Interval) — Apple Maggot, Codling Moth, Eastern Tent Caterpillar, European Apple Sawfly, Eye-Spotted Bud Moth, Flea Beetle, Flea-Twig Leafroller, Green Apple Aphid, Japanese Beetle, Lecanium Scale, Plum Curculio, Redbanded Leafroller, Rose Apple Aphid, San Jose Scale — 70 to 120 lbs. per acre. For Codling Moth and Plum Curculio, spray at petal fall and every 10 to 14 days. Inadequate control is achieved. Orange Tortrix — 50 to 160 lbs. per acre.

3

TURNIPS (14-Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

POULTRY (Chickens, Ducks, Game Birds, Geese, Partridges, Pheasants, Pigeons, Turkey) — Chicken Nit, Fleas, Lice, Northern Fly — 1 lb. per 100 birds. Direct application. Apply by shaker can or hand sprayer. Direct spray to vent and fluff areas. Also treat head and neck for maggot larvae. Repeat if needed. This is a supplement to premise and litter treatments for chicken mites and lice.

SMALL TRAILHOUSES — See Bug, Chicken Nit, Flea, Northern Fly, Mite — 25 lbs per 1000 square feet. Apply to floors, litter, roosts, and adjacent areas with a broom, hand or power duster. For dusting cracks and crevices.

DOG - KENNELS, SLEEPING QUARTERS — 1% to 2% lbs. per 2000 square feet; for ticks, 2% to 5 lbs. per 5000 square feet or less. Apply liberally by gauze-type duster. Forces into cracks and crevices. Repeat as needed. Do not expose kittens or puppies less than 4 weeks of age to treated areas. Do not use on pregnant dogs.

ORNAMENTALS

WOODY SHRUBS, TREES, AND VINES — Sevin injures Boston Ivy and Virginia Creeper. Injury to tender foliage may occur if plants are wet when treated or in the presence of high humidity. Apple Aphid, Bagworm, Birch Leafminers, Boxelder Bug, Boxwood Leaf Miner, Elm Leaf Aphid, Elm Leaf Beetle, Green Moth, Japanese Beetle, Leafhopper, Leafhopper, Bald-faced Hornet, Orange Tortrix, Persimmon Cockroach, Rose Aphid, Rose Slug, Scale, Thrip (exposed), Twospot Spider, Willow Leafhopper — 20 to 40 lbs. per acre. Plant Bug, Psylla — 30 to 40 lbs. per acre.

HERBACEOUS PLANTS AND BULBS (Annual Plants — Herbaceous Plants) — Sevin injures Boston Ivy and Virginia Creeper. Injury to tender foliage may occur if plants are wet when treated or in the presence of high humidity. Blister beetle occurs plants are wet when treated or in the presence of high humidity. Blister Beetle, Flea Beetle, Japanese Beetle, Japanese Leafhopper, Leafhopper, Plant Bug, Psylla, Rosa Aphid, Thrip (exposed) — 30 to 40 lbs. per acre.

23

SQUASH (14-Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

SQUASH (No Pre-Harvest Interval) — Sevin injures watermelons in Florida. Fleaworm, Melonworm — 10 lbs. per acre. Cucumber Beetle, Flea Beetle, Leafhopper, Squash Bug — 20 lbs. per acre.

STRAWBERRIES (1-Day Pre-Harvest Interval) — Meadow Buttercup, Omphacis Leaf Miner, Strawberry Leafroller, Strawberry Weevil — 20 to 40 lbs. per acre.

SUGAR BEETS (14-Day Pre-Harvest Interval) — Grasshopper — 10 to 20 lbs. per acre. Use 10 to 20 lbs. rate on young plants and for controlling nymphs.

SWISS CHARD (14-Day Pre-Harvest Interval) — Flea Beetle, Harpalus Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the principal problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

TOBACCO (In Pure Bed) — Tobacco Flea Beetle — 1 to 2 lbs. per 100 square yards. Apply just before transplanting.

TOBACCO (In Fields) — Cotton Bollworm, Tobacco Flea Beetle — 2 to 2 lbs. per acre. This rate is limited to hand applications directed to the bud and top leaves only. Avoid heavy application as wet, dark leaves & plant injury may result. Leaves dry. Avoid heavy application as wet, dark leaves & plant injury may result. Flea Beetle, Hornworm, Japanese Beetle, June Beetle, Tobacco Budworm — 15 to 25 lbs. per acre. Use full coverage. Aphid, Grasshopper — 30 to 50 lbs. per acre.

TOMATOES (No Pre-Harvest Interval) — Cutworm, Potato Beetle, Flea Beetle, Leafhopper — 10 to 20 lbs. per acre. European Corn Borer, Fall Armyworm, Leafhopper, Flea, Potato Rootworm, Tomato Hornworm, Tomato Rootworm, Tarnished Plant Bug — 20 to 40 lbs. per acre. Climbing cutworm — 20 lbs. per acre.

PEAS (No Pre-Harvest Interval) — Alfalfa Looper, Colorado Potato Beetle, Leafhopper — 20 lbs. per acre. Claims for Colorado Potato Beetle are limited to control of weeds in pea fields to avoid insect contamination in processed beans. Armyworms, Grasshopper — 20 to 40 lbs. per acre.

PEPPERS (No Pre-Harvest Interval) — Colorado Potato Beetle, Leafhopper — 10-20 lbs. per acre. European Corn Borer, Fall Armyworm, Stick Bug, Tomato Hornworm, Tomato Hornworm, Tarnished Plant Bug — 20 to 40 lbs. per acre. Climbing Cutworm — 20 lbs. per acre.

PLUMS (1-Day Pre-Harvest Interval) — Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Tussock Moth — 80 to 120 lbs. per acre. For Codling Moth, apply at petal fall and every 10 to 14 days thereafter until control is achieved. Black Cherry Aphid, Brown Soft Scale, Cherry Anthony, Cherry Fruitworm, Eastern Spotted Bud Moth, Forest Scale, Fruit-Tear Leafroller, Japanese Beetle, Lecanium Scale, Mealy Plum Aphid, Oystershell Scale, Peach Twig Borer, Plum Curculio, Plum Leafhopper, Redheaded Leafroller, San Jose Scale — 20 to 100 lbs. per acre. For optimum scale control apply when crawlers are present. For Plum Curculio, apply at petal fall and every 10 to 14 days thereafter until control is achieved. For optimum scale control apply when crawlers are present.

POTATOES (1-Day Pre-Harvest Interval) — Colorado Potato Beetle, Leafhopper — 10 to 20 lbs. per acre. European Corn Borer, Fall Armyworm, Stick Bug, Tomato Hornworm, Tomato Hornworm, Tarnished Plant Bug — 20 to 40 lbs. per acre.

PRUNES (1-Day Pre-Harvest Interval) — Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Tussock Moth — 80 to 120 lbs. per acre. For Codling Moth, apply at petal fall and every 10 to 14 days thereafter until control is achieved. Black Cherry Aphid, Brown Soft Scale, Cherry Anthony, Cherry Fruitworm, Eastern Spotted Bud Moth, Forest Scale, Fruit-Tear Leafroller, Japanese Beetle, Lecanium Scale, Mealy Plum Aphid, Oystershell Scale, Peach Twig Borer, Plum Curculio, Plum Leafhopper, Redheaded Leafroller, San Jose Scale — 20 to 100 lbs. per acre. For optimum scale control apply when crawlers are present. For Plum Curculio, apply at petal fall and every 10 to 14 days thereafter until control is achieved. For optimum scale control apply when crawlers are present.

• PUMPKINS (No Pre-Harvest Interval) — Scouting injuries, water melons in Florida, Pickleworm, Mexican bean — 12 lbs. per acre. Cucumber Beetle, Flea Beetle, Leafhopper, Squash Bug — 20 lbs. per acre.

10

LAWNS AND TURF — Ants, Carpet Bugs, Cicadas, Cutworms, Earwigs, Fall Armyworms, Flies, Leafhoppers, Millipedes, Mosquitoes, Red Webworms, Ticks — 10 to 20 lbs. per 5,000 square feet. For best results, mow lawn, remove clippings, and apply immediately after rain or watering. Water following treatment. Keep children and pets off lawn material as watered in and grass is dry.

NONAGRICULTURAL LANDS, PASTURE, RANGELAND — Mosquitoes — 5 to 12 lbs. per acre. For oil floor spreader application only. Thorough coverage necessary to effective control. For optimum control, apply in early morning or evening when insects are calm and adults are active. Stay off shrubs and crops. Do not use in 300' where these are important resources.

WOODED AREAS — House Fly, Mosquito, Sandfly — 8 lbs. per acre. Apply with power duster or large hand duster.

DOMESTIC DWELLINGS — 10 to 15 lbs. per 5,000 square feet. Apply to soil surface, mulch, and debris around moist, shady places. Fleas — 5 to 10 lbs. per 5,000 square feet. Apply liberally to parks, yards, lawns and under buildings around the home. Force the dust into cracks and crevices. Repeat as needed. Millipedes — 10 to 15 lbs. per 5,000 square feet. Apply to soil surface, mulch and debris areas and moist, shady places. Mosquitoes — 8 to 10 lbs. per 5,000 square feet. Apply to lawns, shrubbery and other vegetation around the home. Rattlesnakes — 10 lbs. Ticks — 2 1/2 to 5 lbs. per 6,000 square feet. Apply liberally to moist, damp, and vegetation at a height of about 2 feet on lawns, vacant lots, and wooded areas around the home. Repeat if needed. Do not use this product on commercial food areas or food handling establishments, restaurants, or other places where food is prepared or served.

URBAN AND RURAL OUTDOOR AREAS (PUBLIC HEALTH) — Plague Reservoir Host Animals — For use only by and under the supervision of State or Federal Public Health or Wildlife Officials. Fleas — 8 oz. per gallon. Apply by hand or power sprayer. Fleas — 4 oz. per gallon station. Use 4 oz. of bait stations per acre. Fleas — 20 lbs. per acre. Apply by hand or power sprayer as an area spray treatment. Consult your State Fish and Game Agency before applying this product.

• TO AVOID POSSIBLE INJURY TO TENDER FOLIAGE, DO NOT APPLY WHEN FOLIAGE IS WET OR WHEN RAIN OR EXCESSIVE HUMIDITY IS EXPECTED DURING THE NEXT TWO DAYS.

10

RADISHES (9-Day Pre-Harvest Interval) — Flea Beetle, Harvester Bug — 10 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Rootworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug — 20 to 40 lbs. per acre. Where the Cabbage Looper is the primary problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

RASPBERRIES (7-Day Pre-harvest Interval) — Japanese Beetle, Leafhopper, Leaf Roller, Raspberry Aphid — 20 lbs. per acre. Climbing Cutworm, Orange Tortrix, Sawfly, Snowy Tree Cricket — 40 lbs. per acre.

REEDS (10-Day Pre-Harvest Interval) — Grasshoppers — 10 to 20 lbs. per acre. Use 10 to 20 lbs. rate on young plants and/or controlling nymphs. Armyworm, Stick Bug — 20 to 30 lbs. per acre. Use restricted to Mississippi Delta and Texas. To avoid plant injury, Do not spray before heading. A propiconazole oil will be applied. Do not use on the fields in which soybean and/or cotton planting is increased in the cultural practice.

ROTENONE (1-Day Pre-Harvest Interval) — Flea Beetle, Harlequin Bug — 1 to 20 lbs. per acre. Armyworm, Cabbage Looper, Corn Earworm, Diamondback Moth, Imported Cabbageworm, Stick Bug, Tarnished Plant Bug, — 20 to 40 lbs. per acre. Where the Cabbage Looper is the primary problem, use an alternative pesticide specifically recommended for it. For control of small Cabbage Loopers after edible parts start to form, apply at 5 to 7 day intervals.

SALALY (3-Day Pre-Harvest Interval, roots) — Flea Beetle, Leafhopper — 10 to 20 lbs. per acre. Armyworm, Stick Bug — 20 to 40 lbs. per acre.

SORGHUM (Grain, Millet) (1-Day Pre-Harvest Interval, Grain) — No Pre-Harvest Interval for Flea Beetles — 10 to 20 lbs. per acre. Use 10 to 20 lbs. rate on young plants and for controlling nymphs. Armyworm, Corn Earworm, Stick Bug, Thrip, Webworm — 20 to 40 lbs. per acre. Panoplites omnivorus: on grain, straight, direct into the forming heads. Sorghum Nigra — 20 to 2 lbs. per acre. Treat 2 to 4 days after heads have emerged from coat. Direct treatment into forming heads.

• SOYBEANS (No Pre-Harvest Interval) — Flea Beetles, Mexican Bean Beetles — 10 to 20 lbs. per acre. Grasshoppers — 10 to 30 lbs. per acre. Use 10 to 2 lbs. rate for controlling nymphs. Flea beetles as needed. Alfalfa Beetles, Leaf, Southern Beetles, Green Cloverworm, Japanese Beetle, Leafhopper, Flea, Grasshopper, Alfalfa Hopper, Thrip, Velvetbean Capsid — 20 lbs. per acre. Armyworm, Corn Earworm, Stick Bug, Webworm — 20 to 30 lbs. per acre.

11

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants: (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purpose set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests, effectiveness, efficiency to laboratory animals and to plants, and residues on food crops, and user reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTOMATICALLY, ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCL. USES REMEDY AND MANUFACTURER'S SOLE & EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, EXPENSES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT IN RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.